

The Needles in the Haystack survey: searching for X-ray selected normal galaxies

A. E. Georgakakis¹ I. Georgantopoulos¹, S. Basilakos¹, M. Plionis^{1,2}

¹ *Institute of Astronomy & Astrophysics, National Observatory of Athens, I. Metaxa & V. Pavlou, Athens, 15236, Greece*

² *Instituto Nacional de Astrofísica Óptica y Electrónica, AP 51 y 216, 72000, Puebla, Pue, México*

2 February 2008

ABSTRACT

In this paper we present the first results from an on-going serendipitous survey aiming to identify X-ray selected ‘normal’ galaxies (i.e. not AGN dominated) by combining archival XMM-*Newton* data with the Sloan Digital Sky Survey. In the first 4.5 deg² of this program we have identified a total of 11 ‘normal’ galaxy candidates (8 of them with optical spectroscopy) with fluxes $f_X(0.5 - 8 \text{ keV}) \approx 10^{-15} - 10^{-13} \text{ erg s}^{-1} \text{ cm}^{-2}$. These sources are selected to have low X-ray-to-optical flux ratio ($\log f_X/f_{opt} \lesssim -2$), soft X-ray spectral properties and optical spectra, when available, consistent with the presence of a stellar ionising continuum. These sources comprise both early and late type systems at redshifts $z \lesssim 0.2$ with luminosities $L_X(0.5 - 8 \text{ keV}) \approx 10^{39} - 10^{42} \text{ erg s}^{-1}$. This dataset provides the first tight constraint on the surface density of X-ray selected ‘normal’ galaxies at relatively bright fluxes spanning two orders of magnitude ($10^{-15} - 10^{-13} \text{ erg s}^{-1} \text{ cm}^{-2}$). The slope of the ‘normal’ galaxy $\log N - \log S$ in the above flux range is estimated -1.4 ± 0.3 consistent with the euclidean prediction. We also discuss the prospects of ‘normal’ galaxy studies at X-ray wavelengths using both our continuously expanding survey and future X-ray missions.

Key words: Surveys – X-rays: galaxies – X-rays: general

1 INTRODUCTION

In the local Universe ($\lesssim 100 \text{ Mpc}$) ‘normal’ galaxies with X-ray emission dominated by stellar processes have been extensively studied using data from targeted observations of selected systems (e.g. Read, Ponman & Strickland 1997; Shapley, Fabbiano & Eskridge 2001; Zezas et al. 2003; Fabbiano et al. 2003; Jeltema et al. 2003). Although these individual observations provide valuable information on the nature and the X-ray properties of ‘normal’ galaxies they cannot be used for statistical studies (e.g. $\log N - \log S$, luminosity function, clustering properties).

It is only recently that ultra-deep *Chandra* surveys reaching fluxes $f(0.5 - 2 \text{ keV}) \approx 3 \times 10^{-17} \text{ erg s}^{-1} \text{ cm}^{-2}$ (e.g. Brandt et al. 2001; Giacconi et al. 2002; Alexander et al. 2003) have provided the first X-ray selected sample of ‘normal’ galaxy candidates (Hornschemeier et al. 2003; Norman et al. 2004). These objects although too faint to significantly contribute to the X-ray background (XRB) are detected in increasing numbers with decreasing flux and are likely to outnumber powerful AGNs below $f(0.5 - 2 \text{ keV}) \approx 10^{-17} \text{ erg s}^{-1} \text{ cm}^{-2}$ (Hornschemeier et al. 2003).

The studies above however, concentrate on ‘normal’ galaxies at faint X-ray limits ($\approx 10^{-16} \text{ erg s}^{-1} \text{ cm}^{-2}$) and

moderate redshifts $z = 0.1 - 1$ (median ≈ 0.3). At brighter fluxes ($\approx 10^{-15} - 10^{-13} \text{ erg s}^{-1} \text{ cm}^{-2}$) probing, on average, lower redshifts, $z \lesssim 0.2$, our knowledge on these systems remains scanty. This is because ‘normal’ galaxies at bright X-ray fluxes have low sky density. Therefore, wide area surveys are required to compile large enough samples to explore their statistical properties (luminosity function, surface density; Georgakakis et al. 2004). Such a low- z study is essential to interpret the X-ray ‘normal’ galaxies in deeper X-ray surveys and to investigate their evolution. This is particularly important since the X-ray emission is the only observable window available that allows one to probe the evolution of the X-ray binaries and the hot gas in galaxies. This provides information on the star-formation history of the Universe that is complementary to that obtained from other wavelengths (Ghosh & White 2001; Hornschemeier et al. 2002; Georgakakis et al. 2003; Norman et al. 2004).

Wide areal coverage is difficult to achieve with the *Chandra* observatory due its limited field-of-view. On the contrary, XMM-*Newton* with 4 times larger field-of-view provides an ideal platform for such a study. In this paper we exploit the capabilities and the large volume of archival data of the XMM-*Newton* to serendipitously identify ‘normal’ galaxy candidates in public fields selected to overlap

with the Sloan Digital Sky Survey (SDSS; York et al. 2000) to exploit the superb and uniform 5-band optical photometry and spectroscopy available in this area. In this paper we report the first results of this survey (Needles in the Haystack Survey; NHS) using a pilot sample covering a total area of $\approx 4.5 \text{ deg}^2$ in the flux range $f_X(0.5 - 8 \text{ keV}) \approx 10^{-15} - 10^{-13} \text{ erg s}^{-1} \text{ cm}^{-2}$. This dataset is continuously expanded as more SDSS fields and XMM-*Newton* archival data become available. Throughout this paper we adopt $H_0 = 65 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0.3$ and $\Omega_\Lambda = 0.7$.

2 THE SAMPLE

In this paper we use XMM-*Newton* archival observations, with a proprietary period that expired before September 2003, that overlap with first data release of the SDSS (DR1; Stoughton et al. 2002). Only observations that use the EPIC (European Photon Imaging Camera; Strüder et al. 2001; Turner et al. 2001) cameras as the prime instrument operated in full frame mode were employed. For fields observed more than once with the XMM-*Newton* we use the deeper of the multiple observations. A total of 20 fields are used with Galactic N_H in the range $1.4 - 13.2 \times 10^{20} \text{ cm}^{-2}$ (mean $3.7 \times 10^{20} \text{ cm}^{-2}$) and PN good time intervals between 2 and 67 ks (mean 18.5 ks). These observations are complemented with 8 XMM-*Newton* pointings from the northern region of the wide area, shallow XMM-*Newton*/2dF survey that also overlaps with the SDSS (Georgakakis et al. 2004).

The X-ray data reduction, source detection and flux estimation are carried out using methods described in Georgakakis et al. (2004). Source extraction is performed in the 0.5-8 keV merged PN+MOS images using the EWAVELET task of SAS with a detection threshold of 5σ . The extracted sources for each field were visually inspected and spurious detections clearly associated with CCD gaps, hot pixels or lying close to the edge of the field of view were removed. Figure 1 shows the solid angle covered by our survey as a function of the 0.5-8 keV limiting flux.

To optically identify the X-ray sources we follow the method of Downes et al. (1986) as described in Georgakakis et al. (2004) to calculate the probability, P , a given candidate is the true identification. Here we apply an upper limit in the search radius, $r < 7 \text{ arcsec}$ and a cutoff in the probability, $P < 0.05$, to limit the optical identifications to those candidates that are least likely to be chance coincidences.

‘Normal’ galaxy candidates are selected to have (i) extended optical light profile (i.e. resolved) to avoid contamination of the sample by Galactic stars, (ii) X-ray-to-optical flux ratio $\log f_X/f_{opt} < -2$, two orders of magnitude lower than typical AGNs (for the definition of $\log f_X/f_{opt}$ see Georgakakis et al. 2004). The sample of ‘normal’ galaxy candidates is presented in Table 1.

The archival X-ray data used here include targeted observations of nearby normal galaxies with low X-ray-to-optical flux ratio. Such sources have been excluded from Table 1. Moreover, a number of ‘normal’ galaxy candidates although not the prime target of the XMM-*Newton* pointing lie at the same redshift as the prime target and are therefore most likely directly associated with it (e.g. cluster or group members). These sources are marked in Table 1 and are excluded from any statistical analysis.




Figure 1. Solid angle as a function of limiting flux density (5σ) in the total 0.5-8 keV band for our survey.

3 OPTICAL AND X-RAY PROPERTIES

Low luminosity Seyferts also lie in the $\log f_X/f_{opt} \lesssim -1$ regime and may contaminate our ‘normal’ galaxy sample (Lehmann et al. 2001). We explore this possibility using both the X-ray and the optical spectroscopic information available for our sources.

For sources #2, 4, 6, 12 and 15 in Table 1 we can constrain the nature of the central engine using the diagnostic diagrams proposed by Veilleux & Osterbrock (1987) that employ intensity line ratios that are least sensitive to dust reddening ($H\beta/[O III] 5007$, $[S II] 6716+31/H\alpha$, $[O I] 6300/H\alpha$ and $[N II] 6583/H\alpha$). We adopt the classification scheme of Ho, Filippenko & Sargent (1997) and find that sources #2, 6, 12 and 15 have HII type spectra while #4 is classified as a Seyfert 2. As will be discussed below the classification of source #4 is consistent with its absorbed X-ray spectrum.

For the remaining sources in Table 1 the classification using diagnostic diagrams cannot be applied because either they have absorption lines only (#9, 10, 16, 17), no spectroscopic information (#1, 5, 7, 11, 14), or both the $H\beta$ and the $[O III] 5007 \text{ \AA}$ lines, essential for spectral classification, are absent from the optical spectrum (#3, 8, 13). The latter sources have strong absorption features from a dominant evolved stellar population suggesting early and intermediate Hubble types. Galaxies with both absorption and emission lines have been extensively studied by Dressler et al. (1999) and Poggianti et al. (2000) on the basis of the equivalent widths of the $H\delta$ (absorption) and $[O II] 3727 \text{ \AA}$ (emission) lines. Our sources have $EW \approx -1$ (absorption) and $\approx 10 \text{ \AA}$ (emission) for the $H\delta$ and $[O II] 3727 \text{ \AA}$ lines respectively. Following the classification of Dressler et al. (1999) the evidence above suggests e(c)-type galaxies which are believed to be systems with constant star-formation rate over a period of time resulting in a mix of young and evolved stellar populations. For these sources however, we cannot exclude the




Figure 2. Cumulative ‘normal’ galaxy counts. Filled circles are the ‘normal’ galaxy candidates in the present study. We plot 10 instead of 11 points because sources #10 and 15 have the same flux and are plotted as one point. The open circles are the source counts of Hornschemeier et al. (2003). The continuous line is the best fit to the $\log N - \log S$ at bright fluxes from the present sample. The dashed line is the model prediction of the Georgantopoulos et al. (1999) X-ray luminosity function of H II galaxies assuming evolution of the form $L_X \sim (1+z)^3$.

possibility of low-luminosity AGN activity diluted by the evolved stellar population (e.g. Severgnini et al. 2003)

We can further constrain the nature of our ‘normal’ galaxy candidates via X-ray spectral analysis using the XSPEC v11.2 package. We use the C-statistic technique (Cash 1979) specifically developed to extract information from even low signal-to-noise ratio spectra. The data are grouped to have at least one count per bin. An absorbed power-law model is fit to the data fixing the power law index to $\Gamma = 1.9$ and leaving N_H free parameter. The results are shown in Table 1. The majority of the ‘normal’ galaxy candidates have soft X-ray spectral properties suggesting obscuration consistent with the Galactic N_H . The only exception is source #4 which is significantly absorbed.

The final sample of ‘normal’ galaxy candidates that are not associated with the prime target of a given XMM-Newton pointing and not showing evidence (e.g. X-ray or optical spectra) for obscured AGN activity comprises 11 sources, #1, 2, 3, 5, 6, 8, 9, 10, 11, 13, 15 in Table 1. Of these 11 sources only sources #1, 5 and 11 do not have optical spectroscopy available.

4 DISCUSSION

In this paper we report on an on-going program aiming to identify X-ray selected ‘normal’ galaxies in the flux range $f(0.5 - 8 \text{ keV}) \approx 10^{-15} - 10^{-13} \text{ erg s}^{-1} \text{ cm}^{-2}$ by combining archival XMM-Newton data with the SDSS. At present our on-going survey covers an area of 4.5 deg^2 and has identified a total of 11 ‘normal’ galaxy candidates with $\log f_X/f_{opt} < -2$. Unlike deeper samples our sources are rel-




Figure 3. $L_X(0.5 - 8 \text{ keV})$ against redshift. Filled circles are for our sample of ‘normal’ galaxy candidates (including sources associated with the prime target of a given XMM-Newton pointing). Open circles represent the sample of Hornschemeier et al. (2003). The two surveys cover different but complementary regions of the $L_X - z$ space.

atively bright at both the optical and the X-ray wavelengths allowing detailed study of their nature thus, minimising any AGN contamination. Both the soft X-ray spectra and the optical spectroscopic properties are consistent with the absence of nuclear activity.

Figure 2 plots the $\log N - \log S$ of our sample in comparison with that of ‘normal’ galaxy candidates identified in the *Chandra* Deep Field North survey probing fluxes $\approx 10^{-16} \text{ erg s}^{-1} \text{ cm}^{-2}$ (Hornschemeier et al. 2003). The present study provides the only constraints on the number density of ‘normal’ galaxies at relatively bright fluxes in the range $f(0.5 - 8 \text{ keV}) \approx 10^{-15} - 10^{-13} \text{ erg s}^{-1} \text{ cm}^{-2}$ spanning two orders of magnitude. The differential ‘normal’ galaxy counts in the 0.5-8 keV band are fit by a power law yielding a slope of -1.4 ± 0.3 for the cumulative number counts. This is in excellent agreement with the results of Hornschemeier et al. (2003) at fainter fluxes ($-1.46^{+0.28}_{-0.30}$). Figure 2 also shows that our wide area medium-deep serendipitous survey is complementary to the ultra-deep pencil beam *Chandra* observations. This is further demonstrated in Figure 3 plotting the $L - z$ plane for both the present sample and that of Hornschemeier et al. (2003). The two surveys cover different but complementary regions of the parameter space with our sample probing the low- z , high- L_X regime. Such a wide $L - z$ plane coverage is essential to explore the evolution of ‘normal’ galaxies at X-ray wavelengths. Although our sample is not large enough yet for such a study, we plan to expand it to $\approx 20 \text{ deg}^2$ by exploiting the 2nd data release of the SDSS. Our enlarged sample will comprise ≈ 40 ‘normal’ galaxies sufficient for luminosity function estimates providing an anchor point at low redshifts, essential for comparison with higher- z studies (e.g. Hornschemeier et al. 2003; Georgakakis et al. 2003; Norman et al. 2004).

However, even the deepest X-ray samples todate find

ID	α_X (J2000)	δ_X (J2000)	P (per cent)	r (mag)	z	net counts	$f_X(0.5 - 8 \text{ keV})$ ($10^{-14} \text{ erg s}^{-1} \text{ cm}^{-2}$)	$L_X(0.5 - 8 \text{ keV})$ ($10^{41} \text{ erg s}^{-1}$)	N_H (10^{22} cm^{-2})	type ^a
1 ^b	03 58 05.22	+01 09 48.96	0.01	14.93	0.074	53	1.52 ± 0.34	2.36	< 0.04	early
2	08 30 59.21	+52 37 47.75	0.80	17.69	0.1363	48	0.34 ± 0.13	2.01	< 0.13	NL
3	08 31 14.54	+52 42 24.77	0.01	15.24	0.0641	120	0.56 ± 0.10	0.66	< 0.02	EA
4	08 31 39.09	+52 42 05.51	0.02	15.70	0.0587	337	2.07 ± 0.14	2.00	$14.9^{+0.9}_{-2.6}$	NL
5 ^b	08 32 02.58	+52 47 13.74	0.01	16.01	0.105	55	0.25 ± 0.09	0.81	< 0.1	late
6	08 32 28.14	+52 36 22.36	< 0.01	13.82	0.0170	716	6.43 ± 0.29	0.52	< 0.02	NL
7 ^{b,c}	09 17 58.59	+51 51 04.75	0.31	16.66	0.219	22	0.61 ± 0.26	9.76	< 0.06	early
8	09 35 18.39	+61 28 31.58	0.27	16.50	0.1241	54	0.96 ± 0.25	4.48	< 0.08	EA
9	09 36 19.46	+61 27 23.11	0.06	16.55	0.1311	67	0.95 ± 0.21	5.25	< 0.03	AB
10	12 31 47.21	+64 14 01.68	< 0.01	13.01	0.009	67	1.95 ± 0.14	0.047	< 0.02	AB
11 ^b	12 32 53.15	+64 08 50.93	0.17	15.43	0.139	50	1.03 ± 0.26	6.12	< 0.06	late
12 ^c	12 45 31.99	-00 32 08.59	0.01	13.12	0.0055	353	2.83 ± 0.19	0.02	$0.11^{+0.07}_{-0.03}$	NL
13	12 44 52.26	-00 25 49.66	0.01	15.34	0.0823	95	0.57 ± 0.12	1.10	< 0.03	EA
14 ^{b,c}	13 03 01.13	+67 25 17.76	0.16	16.16	0.109	49	1.38 ± 0.35	4.90	$0.01^{+0.01}_{-0.07}$	early
15	13 41 37.85	-00 25 55.30	0.05	16.86	0.053	30	1.95 ± 0.52	1.44	< 0.09	NL
16 ^c	23 53 40.52	-10 24 19.91	0.02	15.07	0.0741	52	2.52 ± 0.52	3.92	< 0.03	AB
17 ^c	23 54 05.69	-10 18 30.64	0.06	15.70	0.0734	50	1.06 ± 0.33	1.61	< 0.02	AB

^a AB: absorption lines; NL: Narrow emission lines; EA: both narrow emission and absorption lines

^b photometric redshift and classification from the SDSS

^c source at the same redshift as the target of the XMM-Newton pointing.

The columns are: 1: identification number; 2: right ascension of the X-ray source; 3: declination of the X-ray source;

4: probability the optical counterpart is a spurious alignment; 5: optical magnitude; 6: spectroscopic or photometric redshift from the SDSS;

7: 0.5-8 keV net counts; 8: X-ray flux in the 0.5-8 keV spectral band in units of $10^{-14} \text{ erg s}^{-1} \text{ cm}^{-2}$;

9: 0.5-8 keV X-ray luminosity in units of $10^{41} \text{ erg s}^{-1}$; 10: N_H from the X-ray spectrum assuming $\Gamma = 1.9$; 11: spectral type.

Table 1. The candidate ‘normal’ galaxy sample.

‘normal’ galaxies at relatively low- z (median redshift $z \approx 0.3$; Hornschemeier et al. 2003) significantly lower than the star-formation peak at $z \approx 1 - 2$. This underlines the need for deeper X-ray observations to use X-ray selected ‘normal’ galaxies as cosmological probes. The ESA *XEUS* mission with its unparalleled sensitivity will provide a great leap forward in these studies. Assuming the goal resolution of about 2 arcsec FWHM for its mirrors, *XEUS* will attain, in just 20 ks, about equal sensitivity to the deepest (2 Ms) *Chandra* exposure. Extrapolation of the $\log N - \log S$ relation derived here to $f_X(0.5 - 8) = 10^{-16} \text{ erg cm}^{-2} \text{ s}^{-1}$ (i.e. approximate flux limit of the 2 Ms *Chandra* Deep Field North; Alexander et al. 2003), suggests that about 10 galaxies will be detected in the 25 arcmin² *XEUS* Wide Field Imager. Therefore, a few tens of typical exposures with *XEUS* will readily accumulate a large sample of X-ray selected galaxies for the study of the galaxy luminosity function and evolution at redshifts $z \lesssim 1$ with a median redshift of $z \sim 0.3$. Deeper exposures are necessary to probe the star-formation peak at higher redshifts. In a deep 200 ks exposure, *XEUS* is expected to probe fluxes as faint as $f_X(0.5 - 8 \text{ keV}) \approx 5 \times 10^{-18} \text{ erg s}^{-1} \text{ cm}^{-2}$ and to detect more than 500 galaxies (an estimate obtained by extrapolating our $\log N - \log S$). At these flux levels the mean redshifts probed are $z \sim 1$ as derived from the X-ray luminosity function of Georgantopoulos et al. (1999) assuming evolution of the form $L_X \sim (1+z)^3$. Therefore, combination of the local XMM-Newton galaxy samples with the *XEUS* data will allow an unprecedented study of the star-formation history at X-ray wavelengths.

REFERENCES

- Alexander D., et al., 2003, *AJ*, 126, 539
Brandt W. N., et al., 2001, *AJ*, 122, 1.
Cash W., 1979, *ApJ*, 228, 939
Downes A. J. B., Peacock J. A., Savage A., Carrie D. R., 1986, *MNRAS*, 218, 31
Dressler A., Smail I., Poggianti B. M., Butcher H., Couch W. J., Ellis R. S., Oemler A. Jr., 1999, *ApJS*, 122, 51
Fabbiano G., Krauss M., Zezas A., Rots A., Neff S., 2003, *ApJ*, 598, 272
Georgakakis A., et al., 2004, *MNRAS*, 349, 135
Georgakakis A., Hopkins A. M., Sullivan M., Afonso J., Georgantopoulos I., Mobasher B., Cram L. E., 2003, *MNRAS*, 345, 939
Georgantopoulos I., Basilakos S., Plionis M., 1999, *MNRAS*, 305, L31.
Ghosh P., White N. E., 2001, *ApJ*, 559, 97.
Giacconi R., et al., 2002, *ApJS*, 139, 369
Ho L. C., Filippenko A. V., Sargent W., 1997, *ApJS*, 112, 315.
Hornschemeier A. E., Brandt W. N., Alexander D. M., Bauer F. E., Garmire G. P., Schneider D. P., Bautz M. W., Chartas G., 2002, *ApJ*, 568, 82.
Hornschemeier A. E. et al., 2003, *AJ*, 126, 575
Jeltema T. E., Canizares C. R., Buote D. A., Garmire G. P., 2003, *ApJ*, 585, 756
Lehmann I., et al., 2001, *A&A*, 371, 833.
Norman C., et al., 2004, *ApJ*, in press, astro-ph/0402140
Poggianti B. M., Smail I., Dressler A., Couch W. J., Barger A. J., Butcher H., Ellis R. S., Oemler A. Jr., 1999, *ApJ*, 518, 576
Read A. M., Ponman T. J., Strickland D. K., 1997, *MNRAS*, 286, 626.

- Severgnini P. et al., 2003, A&A, 406, 483
Shapley A., Fabbiano G., Eskridge P. B., 2001, ApJS, 137, 139.
Stoughton C., et al., 2002, AJ, 123, 485.
Strüder L., Briel U., Dennerl K., et al. 2001, A&A, 365, L18.
Turner M. J. L., Abbey A., Arnaud M., et al., 2001, A&A, 365, L27.
Veilleux S., Osterbrock D. E., 1987, ApJS, 63, 295
York D. G., et al., 2000, AJ, 120, 1579. 2003ApJ...599L..73Z
Zezas A., Hernquist L., Fabbiano G., Miller J., 2003, ApJ, 599L, 73